



BROADLEAF WEED CONTROL

CONTAINS **TRIMEC®** HERBICIDE

Controls chickweed, dandelion, knotweed, plantain, henbit, spurge and many other weeds as listed.

TURF FERTILIZER

22-0-5

Trimec, PCSCU, 1% Fe GUARANTEED ANALYSIS

Total Nitrogen (N)	22.0%
22.0% Urea nitrogen*	
Soluble Potash (K ₂ O)	5.0%
Iron (Fe)	1.0%

Primary Plant foods derived from: Urea, Muriate of Potash and Poly Coated Sulfur Coated Urea. Iron derived from Iron Oxy-Sulfate.

*6.6% slowly available urea nitrogen from poly coated sulfur coated urea.

544-3030

genericpocket

ACTIVE INGREDIENTS

Dimethylamine salt of 2,4-dichlorophenoxyacetic acid	1.040%
Dimethylamine salt of (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid	0.230%
Dimethylamine salt of dicamba (3,6-dichloro-o-anisic acid)	0.108%

INERT INGREDIENTS	98.622%
TOTAL	100.000%

THIS PRODUCT CONTAINS:

17.20 lbs. 2,4-dichlorophenoxyacetic acid equivalent per ton or 0.86%
3.80 lbs. (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per ton or 0.19%
1.80 lbs. 3,6-dichloro-o-anisic acid equivalent per ton or 0.09%

Isomer Specific by AOAC Methods

TRIMEC® is a registered trademark of PBI/GORDON CORPORATION

EPA REG. NO. 2217-561-57131

EPA EST. NO. 57131-WI-1



KEEP OUT OF REACH OF CHILDREN CAUTION



READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Harmful if swallowed. Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Wear long-sleeved shirt, long pants, socks, shoes and chemical-resistant gloves made of any waterproof material.

USER SAFETY REQUIREMENTS:

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

- Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

FIRST AID	
If swallowed:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Do not give anything by mouth to an unconscious person.
If on skin or clothing:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15 - 20 minutes.• Call a poison control center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment information.	

ENVIRONMENTAL HAZARDS:

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

PRECAUTIONS AND RESTRICTIONS:

- Do not apply this product in a way that will contact workers or other persons, either directly or through drift.
- Only protected handlers may be in the area during application.
- Do not allow people or pets to enter the treated area until dusts have settled.

OTHER USE PRECAUTIONS:

- Do not use this product on or near desirable plants, including within the dripline of the roots of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn or garden, and sweep any product landing on the driveway, sidewalk, gutter, or street, back onto the treated area. To prevent product run-off, do not overwater the treated area to the point of runoff or apply when raining or when rain is expected that day.
- Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

STORAGE & DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING: Nonrefillable container: Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available or dispose of empty bag in a sanitary landfill, or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WHERE TO USE

This product is a Trimec® herbicide containing fertilizer, 2,4-D, mecoprop-p, and dicamba. This granular weed and feed provides plant nutrients and controls a broad spectrum of broadleaf weeds in turfgrass. Supplemental nitrogen, phosphorus, and potash should be applied according to soil tests or State Extension Service recommendations. This product is intended to be applied by commercial applicators, noncommercial users, lawn care operators, and landscape personnel. Cool season and warm season grasses established in institutional, ornamental, and residential sites may be treated. Institutional sites are defined as turf areas around properties or facilities providing a service to public or private organizations; for example: hospitals, nursing homes, schools, museums, libraries, sport facilities (including golf courses), and office buildings may be treated. Ornamental sites include turfgrass established around residences, parks, streets, retail outlets, cemeteries, and institutional buildings. Finally, residential sites are defined as areas associated with the household or home life including, but not limited to apartment complexes, condominiums, and patient care areas of nursing homes, mental institutions, hospitals, or convalescent homes. This product may be applied to newly seeded cool season turfgrass. Applications are suitable only after the third mowing or after the grass seedlings have tillered and have developed an adequate secondary root system. This product may be applied to newly established warm season turfgrass. Treatments should be delayed until 3 to 4 weeks after the sodding, sprigging, or plugging operations, and low application rates (½ of the recommendation for established turfgrass) are suggested. A uniform broadcast distribution of product on the treated area is essential. Do not exceed the specified application rate

for any area. Do not use this product on St. Augustinegrass, dichondra, carpetgrass, closely-mowed bentgrass putting greens, or turf containing desirable clovers.

HOW MUCH TO APPLY

For Kentucky bluegrass, fescue spp., perennial ryegrass, bermudagrass, and zoysiagrass: Apply 174 pounds of this product per acre of established turfgrass. Or the equivalent application rates are 4.0 pounds of product per 1,000 square feet of turfgrass. The maximum broadcast application rate is 4.0 pounds of product per 1,000 sq.ft. per application. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 8.0 pounds of product per 1,000 sq.ft.. Sequential or follow-up applications may be required for dense infestations of perennial and biennial weeds. Also, adverse environmental conditions prior or after application may reduce the effectiveness. If required, the treatment interval between the first and second applications is 30 days. Drop spreaders, rotary spreaders, and other granular equipment should be calibrated for accurate and uniform applications of this product. Calibrate the equipment by applying the product over a measured or known area and adjust the equipment according to the recommendations of the equipment manufacturer.

APPLICATION SUGGESTIONS:

1. Spring and fall applications are suggested for the best control. However, this product can be applied any time broadleaf weeds are present and actively growing.
2. Mow the turfgrass 1 to 2 days before application.
3. Water the turfgrass thoroughly (> ½ inch) at least 1 to 2 days before application.
4. Schedule applications in the morning when dew is present. Or, sprinkle or irrigate lightly (<¼ inch) before application to provide maximum control.
5. Rainfall or irrigation within 24 hours after application may reduce the effectiveness. Do not irrigate or water the turfgrass within 24 hours after application.

CALIBRATION INSTRUCTIONS:

To provide proper distribution, calibrate spreader before application. For 4.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 250 square feet and adjust to apply at the rate of 1 pound of product per 250 square feet.

4 lbs Per 1,000	
SPREADERS	Sq. Ft.
Lesco Rotary	F-G
Spyker 24-22	4
Scott's R8A	1
Prizelawn CB Rotary	4
Earthway 2400 Rotary	17
Lely Broadcast	3.4

DISTRIBUTION AND SPREADER PATTERNS: Always apply with a properly calibrated rotary or drop type fertilizer spreader to distribute the granules evenly. Never apply by hand.

COVERAGE: To determine how much you need to apply, multiply the length of your lawn by its width.

This equals your total square footage. Deduct all non-lawn areas (house, gardens, etc.). One bag covers up to 12,500 square feet.

SINGLE PASS DISTRIBUTION: First spread the borders of your lawn. Then spread the center by spreading in straight line passes. Spread at a uniform walking speed to insure even coverage. To avoid gaps and streaks, first spread the border and fill in center with uniform, parallel passes. Keep this product off sidewalks, patios, driveways, etc. This fertilizer contains iron which will stain concrete and painted surfaces. If it comes in contact with concrete surfaces, sweep off immediately before flushing with water. If concrete is wet when contact occurs, staining may be impossible to prevent. Be careful when using rotary spreaders which scatter fertilizer beyond lawn edge and onto concrete.

WEEDS CONTROLLED:

Dandelion, chickweed, plantain, and clover are controlled effectively with this product. A partial list of susceptible weeds is shown below.

Bedstraw	Chicory	Ground Ivy	Lepedeza	Pigweed	Ragweed	Thistle	Yarrow
Black medic	Clovered	Heallall	Mallow	Plantain	Sheep sorrel	Wild Carrot	
Buckhorn	Clover, white	Henbit	Morningglory	Poison Ivy	Sheepspurse	Wild Garlic	and other
Burdock	Dandelion	Knotweed	Oxalis	Poison Oak	Speedwell	Wild Lettuce	broadleaf weeds
Chickweed, common	Curly Dock	Lambsquarters	Pepperweed	Purslane, common	Spotted Spurge	Wild Onion	

LIMITED WARRANTY AND DISCLAIMER.

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use. TO THE EXTENT ALLOWED BY LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law. To the extent allowed by law, the terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

EC GROW

Distributed and Guaranteed By Eau Claire Co-op Oil Co.
P.O. Box 837, Eau Claire, WI 54602
www.ecgrow.com

Nonrefillable Container
NET WT. 50 lbs. (22.68 kg)

Doc.# RP2217-025-12, Rev. Date: 8/6/12